

CD28 Monoclonal Mouse Antibody (C28/1636)

Product Description

This antibody recognizes a glycoprotein of 44-88 kDa, which is identified as CD28. It is the critical T-cell co-stimulatory receptor which provides to the cell the important second activation signal by binding CD80 and CD86 that are expressed by antigen presenting cells. Besides its co-stimulation role, CD28 functions in preventing T-cells from anergic hyporesponsive state or from undergoing premature apoptotic cell death. CD28 is also expressed on human fetal NK cells and some NK cell lines, whereas on murine NK cells the CD28 expression is much broader.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the <u>CF® Dye Brochure</u> for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <u>order@biotium.com</u> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 1636, Anti-CD28 (C28/1636)

Call us : 800-304-5357 Email: btinfo@biotium.com

Product attributes

Flouder attributes				
Antibody number	#1636			
Antibody reactivity (target)	CD28			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	C28/1636			
Isotype	lgG1			
Molecular weight	44 kDa (monomer); 88 kDa (dimer)			
Synonyms	T-cell-specific surface glycoprotein CD28; Tp44			
Human gene symbol	CD28			
Entrez gene ID	940			
SwissProt	P10747			
Unigene	591629			
Immunogen	Recombinant human full-length CD28 protein			
Antibody target cellular localization	Plasma membrane			
Verified antibody applications	Flow (verified)			
	Human			
Species reactivity	Human			
Species reactivity Antibody application notes	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user			
• •	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mlL, Flow Cytometry 0.5-1 ug/mllc cells/0.1 mL, Optimal dilution for a			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user			
Antibody application notes Positive control	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user Jurkat cells or human lymph nodes and tonsils.			
Antibody application notes Positive control Shipping condition	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user Jurkat cells or human lymph nodes and tonsils. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
Antibody application notes Positive control Shipping condition Storage Conditions	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user Jurkat cells or human lymph nodes and tonsils. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when			
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mIlion cells/0.1 mL, Optimal dilution for a specific application should be determined by user Jurkat cells or human lymph nodes and tonsils. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended			
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user Jurkat cells or human lymph nodes and tonsils. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; D2 mg/mL in PBS/0.05% BSA/0.05% azide, PURIFIEd; D2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; D2 mg/mL in PBS/0.05% BSA/0.05% azide, PURIFIEd; BSA-free: 1 mg/mL in PUS			
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mlL, Flow Cytometry 0.5-1 ug/mlL or els/0.1 mL, Optimal dilution for a specific application should be determined by user Jurkat cells or human lymph nodes and tonsils. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% azide, Purified; 0.2 mg/mL in PBS/0.05% azide, Purified; 0.2 mg/mL in PBS without azide			
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunolluorescence: 0.5-1 ug/mll. Flow Cytometry 0.5-1 ug/mll or els/0.1 mL, Optimal dilution for a specific application should be determined by user Jurkat cells or human lymph nodes and tonsils. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% Saide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% Saide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% saide, Purified, DSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% at a prime priman prime priman prime prime prime prime prime prime pr			

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on May 9, 2025 at 03:57:33 AM. Visit product page to check for updated information before use. Product link: <u>http://54.245.69.9/product/mouse-monoclonal-anti-cd28-c281636/</u>