Chromogranin A Recombinant Monoclonal Rabbit Antibody (CHGA/1815R)



Product Description

Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, paragangliomas, and other neuroendocrine tumors. Co-expression of chromogranin A and neuron specific enolase (NSE) is common in neuroendocrine neoplasms. Reportedly, co-expression of certain keratins and chromogranin indicates neuroendocrine lineage. The presence of strong anti-chromogranin staining and absence of anti-keratin staining should raise the possibility of paraganglioma. The co-expression of chromogranin and NSE is typical of neuroendocrine neoplasms. Most pituitary adenomas and prolactinomas readily express chromogranin.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 1815, Anti-Chromogranin A (CHGA/1815R)

Product attributes

Antibody number	#1815			
Antibody reactivity (target)	CHGA, Chromogranin A			
Antibody type	Primary			
Host species	Rabbit			
Clonality	Recombinant Monoclonal			
Clone	CHGA/1815R			
Isotype	IgG, kappa			
Molecular weight	68-75 kDa			
Synonyms	Beta-Granin; CGA; CHGA; Chromogranin A Parathyroid Secretory Protein 1; ER-37; Pancreastatin; Parastatin; Pituitary Secretory Protein I; SP-I; Vasostatin I or II			
Human gene symbol	CHGA			
Entrez gene ID	1113			
SwissProt	P10645			
Unigene	150793			
Immunogen	Recombinant full-length human CHGA protein			
Verified antibody applications	IHC (FFPE) (verified)			
Antibody target cellular localization	Vesicular			
Species reactivity	Human			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user			
	20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application			
Positive control	20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application			
Positive control Shipping condition	20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user PC12 cells. Adrenal gland, bowel, thyroid, pancreas, or			
	20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user PC12 cells. Adrenal gland, bowel, thyroid, pancreas, or pheochromocytoma.			
Shipping condition	20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user PC12 cells. Adrenal gland, bowel, thyroid, pancreas, or pheochromocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
Shipping condition Storage Conditions	20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Western blotting 0.5-1 ug/ml., Optimal dilution for a specific application should be determined by user PC12 cells. Adrenal gland, bowel, thyroid, pancreas, or pheochromocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when			
Shipping condition Storage Conditions Shelf life	20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Western blotting 0.5-1 ug/ml., Optimal dilution for a specific application should be determined by user PC12 cells. Adrenal gland, bowel, thyroid, pancreas, or pheochromocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user PC12 cells. Adrenal gland, bowel, thyroid, pancreas, or pheochromocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05%			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user PC12 cells. Adrenal gland, bowel, thyroid, pancreas, or pheochromocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation Validated in protein array	20 min, Flow Cytometry 0.5-1 ug/mllion cells/0.1 ml., Western blotting 0.5-1 ug/ml., Optimal dilution for a specific application should be determined by user PC12 cells. Adrenal gland, bowel, thyroid, pancreas, or pheochromocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS without azide Monospecific			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation Validated in protein array Antibody research areas	20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Western blotting 0.5-1 ug/ml., Optimal dilution for a specific application should be determined by user PC12 cells. Adrenal gland, bowel, thyroid, pancreas, or pheochromocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/ml in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/ml. in PBS/0.05% BSA/Purified: 0.2 mg/ml. in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/ml. in PBS/0.05% BSA/0.05% azide, Purified: Denotronlogy, Neuroscience Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in			

Call us: 800-304-5357 Email: btinfo@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on February 12, 2025 at 11:00:48 AM. Visit product page to check for updated information before use. Product link: http://54.245.69.9/product/recombinant-rabbit-monoclonal-anti-chromogranin-a-chga-chga1815r/