GAD1 / GAD67 Monoclonal Mouse Antibody (GAD1/2563)



Product Description

This MAb recognizes a protein of 67 kDa, which is identified as glutamic acid decarboxylase 1 (GDA1). There are two forms of glutamic acid decarboxylases (GADs) that are found in the brain: GAD65 (also known as GAD2) and GAD67 (also known as GAD1. GAD65 and GAD67 are members of the group II decarboxylase family of proteins and are responsible for catalyzing the rate-limiting step in the production of GABA (γ-aminobutyric acid) from L-glutamic acid. Although both GAD s are found in the brain, GAD65 localizes to synaptic vesicle membranes in nerve terminals, while GAD67 is distributed throughout the cell. GAD67 is responsible for the basal levels of GABA synthesis. In the case of a heightened demand for GABA in neurotransmission, GAD65 will transiently activate to assist in GABA production. The loss of GAD65 is detrimental and can impair GABA neurotransmission, however the loss of GAD67 is lethal.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 2563, Anti-GAD1/GAD67 (GAD1/2563)

.

Cell/tissue expression

| Product attributes | | | | |
|---------------------------------------|--|--|--|--|
| Antibody number | #2563 | | | |
| Antibody reactivity (target) | GAD1, GAD67 | | | |
| Antibody type | Primary | | | |
| Host species | Mouse | | | |
| Clonality | Monoclonal | | | |
| Clone | GAD1/2563 | | | |
| Isotype | IgG2b, kappa | | | |
| Molecular weight | 67 kDa | | | |
| Synonyms | 67kDa glutamic acid decarboxylase; CPSQ1; DCE1; GAD67; GAD1; Glutamate decarboxylase 1; SCP | | | |
| Human gene symbol | GAD1 | | | |
| Entrez gene ID | 2571 | | | |
| SwissProt | Q99259 | | | |
| Unigene | 420036 | | | |
| Immunogen | Recombinant human GAD1 (GAD67) protein fragment (around aa 72-135) (exact sequence is proprietary) | | | |
| Verified antibody applications | Flow (intracellular) (verified) | | | |
| Antibody target cellular localization | Cytoplasmic, Plasma membrane, Vesicular | | | |
| Species reactivity | Human | | | |
| Antibody application notes | Higher concentration may be required for direct detection using orimary antibody conjugates than for indirect detection with secondary antibody, ELISA: For coating order antibody without BSA, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at TT, Western: 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, DH 6.0, or 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user | | | |
| Positive control | K-562 or HEK293 Cells. Pancreas | | | |
| Shipping condition | Room temperature | | | |
| Storage Conditions | Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C | | | |
| Shelf life | Guaranteed for at least 24 months from date of receipt when stored as recommended | | | |
| Regulatory status | For research use only (RUO) | | | |
| Antibody/conjugate formulation | Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide | | | |
| Validated in protein array | Monospecific | | | |
| Antibody research areas | Neuroscience | | | |
| Product origin | roduct may contain either bovine serum albumin (BSA) from ovine serum (Bos taurus), or recombinant BSA produced in hinese hamster ovary cells. Inquire for the specific lot. | | | |
| • "" | | | | |

Call us: 800-304-5357 Email: btinfo@biotium.com

| Antibody # prefix | Conjugation | Ex/Em (nm) | Laser line | Detection channel | Dye Features |
|-------------------|-----------------------|------------|------------|-----------------------------|------------------|
| BNC04 | CF®405S | 404/431 | 405 | DAPI (microscopy), AF405 | CF®405S Features |
| BNC88 | CF®488A | 490/515 | 488 | GFP, FITC | CF®488A Features |
| BNC68 | CF®568 | 562/583 | 532, 561 | RFP, TRITC | CF®568 Features |
| BNC94 | CF®594 | 593/614 | 561 | Texas Red® | CF®594 Features |
| BNC40 | CF®640R | 642/662 | 633-640 | Cy®5 | CF®640R Features |
| BNC47 | CF®647 | 650/665 | 633-640 | Cy®5 | CF®647 Features |
| BNC74 | CF®740 | 742/767 | 633-685 | 775/50 | CF®740 Features |
| BNCB | Biotin | N/A | N/A | N/A | |
| BNUB | Purified | N/A | N/A | N/A | |
| BNUM | Purified, BSA-free | N/A | N/A | N/A | |

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on May 9, 2025 at 04:24:37 PM. Visit product page to check for updated information before use. Product link: http://54.245.69.9/product/gad1-gad67-monoclonal-mouse-antibody-gad1-2563/