

OLIG2 Monoclonal Mouse Antibody (OLIG2/2400)



Product Description

Olig2, a basic helix-loop-helix transcription factor, is involved in oligodendroglial specification. Olig2 expression has been reported in most glial tumors, such as oligodendrogliomas and astrocytomas. Although more than half of glioblastomas are positive for Olig2, expression is very weak in terms of both percentage of labeled cells and intensity. No Olig2 expression has been found in the non-glial tumors including neuro-epithelial tumors, ependymomas, sub-ependymomas, medulloblastomas, and non-neuroepithelial tumors, such as CNS lymphomas, meningiomas, schwannomas, atypical teratoid / rhabdoid tumor, and haemangioblastomas. Compared to the strong staining seen in glioma samples, a weak expression is observed in non-tumoral brain tissue (gliosis).

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 2400, Anti-OLIG2 (OLIG2/2400)

Call us : 800-304-5357 Email: btinfo@biotium.com

Product attributes

Product attributes				
Antibody number	#2400			
Antibody reactivity (target)	OLIG2			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	OLIG2/2400			
Isotype	IgG1, kappa			
Molecular weight	30-40 kDa			
Synonyms	Basic helix loop helix protein class B1 (bHLHB1); basic helix-loop-helix protein 19 (bHLHe19); OLIG2; Oligodendrocyte lineage transcription factor 2; Oligodendrocyte specific bHLH transcription factor 2; Oligodendrocyte transcription factor 2; Protein kinase C-binding protein 2 (PRKCBP2); RACK17			
Human gene symbol	OLIG2			
Entrez gene ID	10215			
SwissProt	Q13516			
Unigene	176977			
Immunogen	Recombinant fragment of human OLIG2 protein (around aa 1-141) (exact sequence is proprietary)			
Verified antibody applications	IHC (FFPE) (verified)			
Antibody target cellular localization	Nucleus			
loculization				
Species reactivity	Human			
	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user			
Species reactivity	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal			
Species reactivity Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user			
Species reactivity Antibody application notes Positive control	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user THP-1 cells. Astrocytoma			
Species reactivity Antibody application notes Positive control Shipping condition	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user THP-1 cells. Astrocytoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, optimal dilution for a specific application should be determined by user THP-1 cells. Astrocytoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when			
Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions Shelf life	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user THP-1 cells. Astrocytoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended			
Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin:): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user THP-1 cells. Astrocytoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL			
Species reactivity Antibody application notes Antibody application notes Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin:): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user THP-1 cells. Astrocytoma Room temperature Store at 2 to 8 ° C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% azide, Purified: 0.2 mg/mL in PBS without azide			
Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin:): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user THP-1 cells. Astrocytoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/without azide Monospecific			
Species reactivity Antibody application notes Antibody application notes Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation Validated in protein array Antibody research areas	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user THP-1 cells. Astrocytoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS/0.05% bSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS/0.05% without azide Monospecific Cancer, Neuroscience, Transcription factors Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in			

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on February 12, 2025 at 07:46:06 PM. Visit product page to check for updated information before use. Product link: http://54.245.69.9/product/olig2-monoclonal-mouse-antibody-olig2-2400/