## CD22 / BL-CAM Monoclonal Mouse Antibody (RFB4)



## **Product Description**

Recognizes a protein of 130-140 kDa, identified as CD22 (also known as BL-CAM). CD22 expression is restricted to normal and neoplastic B cells and is absent from other haemopoietic cell types. In B-cell ontogeny, CD22 is first expressed in the cytoplasm of pro-B and pre-B cells, and on the surface as B cells mature to become lgD. It is not expressed by plasma cells, CD22 is found highly expressed in follicular mantle and marginal zone B-cells, and while germinal center B-cells are relatively weak. CD22 is a member of the immunoglobulin superfamily and serves as an adhesion receptor for sialic acid-bearing ligands expressed on erythrocytes and all leukocyte classes. It also associates with tyrosine kinases and play a role in signal transduction and B-cell activation.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <a href="mailto:order@biotium.com">order@biotium.com</a> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 1594, Anti-BL-CAM|CD22 (RFB4)

## Product attributes

Product attributes			
Antibody number	#1594		
Antibody reactivity (target)	BL-CAM, CD22		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	RFB4		
Isotype	IgG1, kappa		
Molecular weight	130-140 kDa		
Synonyms	B-lymphocyte cell adhesion molecule (BL-CAM); B-cell receptor CD22; CD22; Lectin 2; Lyb8; Sialic acid-binding Ig-like lectin 2 (Siglec-2); SIGLEC2; T-cell surface antigen Leu-14		
Human gene symbol	CD22		
Entrez gene ID	933		
SwissProt	P20273		
Unigene	579691		
Immunogen	Human tonsil lymphocytes.		
Verified antibody applications	Flow (verified)		
Antibody target cellular localization	Plasma membrane		
Species reactivity	Human		
	Human Flow, surface (published for clone), Functional studies (published for clone)		
Species reactivity Expected antibody	Flow, surface (published for clone), Functional studies		
Species reactivity Expected antibody applications	Flow, surface (published for clone), Functional studies (published for clone)  ELISA: For coating, order antibody without BSA; Flow cytometry: 0.5-1 ug/million cells; Immunofluorescence: 0.5-1 ug/mL; Optimal dilution for a specific application should be determined., Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with		
Species reactivity Expected antibody applications Antibody application notes	Flow, surface (published for clone), Functional studies (published for clone)  ELISA: For coating, order antibody without BSA; Flow cytometry: 0.5-1 ug/million cells; Immunofluorescence: 0.5-1 ug/mill. Optimal didution for a specific application should be determined., Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody  Ramos, Raji or Daudi cells. Human Tonsil, Lymph Node or		
Species reactivity Expected antibody applications Antibody application notes Positive control	Flow, surface (published for clone), Functional studies (published for clone)  ELISA: For coating, order antibody without BSA; Flow cytometry: 0.5-1 ug/million cells; Immunofluorescence: 0.5-1 ug/mil.; Optimal dilution for a specific application should be determined., Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody  Ramos, Raji or Daudi cells. Human Tonsil, Lymph Node or Spleen.		
Species reactivity Expected antibody applications Antibody application notes  Positive control Shipping condition	Flow, surface (published for clone), Functional studies (published for clone)  ELISA: For coating, order antibody without BSA; Flow cytometry: 0.5-1 ug/million cells; Immunofluorescence: 0.5-1 ug/mil; Optimal dilution for a specific application should be determined, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody  Ramos, Raji or Daudi cells. Human Tonsil, Lymph Node or Spleen.  Room temperature  Note: store BSA-free antibodies at -10 to -35°C, Store at 2 to 8		
Species reactivity  Expected antibody applications  Antibody application notes  Positive control  Shipping condition  Storage Conditions	Flow, surface (published for clone), Functional studies (published for clone)  ELISA: For coating, order antibody without BSA; Flow cytometry: 0.5-1 ug/million cells; Immunofluorescence: 0.5-1 ug/mil; Optimal dilution for a specific application should be determined, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody  Ramos, Raji or Daudi cells. Human Tonsil, Lymph Node or Spleen.  Room temperature  Note: store BSA-free antibodies at -10 to -35°C, Store at 2 to 8°C, Protect fluorescent conjugates from light		
Species reactivity  Expected antibody applications  Antibody application notes  Positive control  Shipping condition  Storage Conditions  Regulatory status  Antibody/conjugate	Flow, surface (published for clone), Functional studies (published for clone)  ELISA: For coating, order antibody without BSA; Flow cytometry: 0.5-1 ug/mll; on cells; Immunofluorescence: 0.5-1 ug/ml.; Optimal dilution for a specific application should be determined., Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody  Ramos, Raji or Daudi cells. Human Tonsil, Lymph Node or Spleen.  Note: store BSA-free antibodies at -10 to -35°C, Store at 2 to 8°C, Protect fluorescent conjugates from light  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/Purified: 0.2 mg/mL in PBS/0.05% BSA/D.05% BSA/O.05% SA/O.05% SA/		
Species reactivity Expected antibody applications Antibody application notes  Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation	Flow, surface (published for clone), Functional studies (published for clone)  ELISA: For coating, order antibody without BSA; Flow cytometry: 0.5-1 ug/mlillon cells; Immunofluorescence: 0.5-1 ug/ml.; Optimal dilution for a specific application should be determined., Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody  Ramos, Raji or Daudi cells. Human Tonsil, Lymph Node or Spleen.  Room temperature  Note: store BSA-free antibodies at -10 to -35°C, Store at 2 to 8°C, Protect fluorescent conjugates from light  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide  Guaranteed for at least 24 months from date of receipt when		
Species reactivity  Expected antibody applications  Antibody application notes  Positive control  Shipping condition  Storage Conditions  Regulatory status  Antibody/conjugate formulation  Shelf life	Flow, surface (published for clone), Functional studies (published for clone)  ELISA: For coating, order antibody without BSA; Flow cytometry: 0.5-1 ug/mlil. 0.5-1 ug/mlil		

Call us: 800-304-5357 Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

## References

Note: References for this clone sold by other suppliers may be listed for expected applications.

- 1. Clin Cancer Res (2005) 11(21): 7920-7928. (Flow, surface; functional studies)
- 2. Biomolecules (2020) 10: 1140. (oncotoxins)