## Ep-CAM / CD326 Monoclonal Mouse Antibody (MOC-31)

## **Product Description**

The epitope of this antibody is located in the first EGF-like repeat domain (EGF1) between amino acids 27-59 of Ep-CAM.EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Catalog number key for antibody number 0380, Anti-CD326|Ep-CAM (MOC-31)

## Call us: 800-304-5357 Email: techsupport@biotium.com

## **Product attributes**

Antibody number	#0380				
Antibody reactivity (target)	CD326, Ep-CAM				
Antibody type	Primary				
Host species	Mouse				
Clonality	Monoclonal				
Clone	MOC-31				
Isotype	IgG1, kappa				
Molecular weight	40-43 kDa				
Synonyms	Adenocarcinoma-associated Antigen; Cell Surface Glycoproteir Trop-1; EGP2; EGP314; EGP40; Epithelial Cell Adhesion Molecule; Epithelial Glycoprotein 314; ESA; KSA; TACD1; TROP1; Tumor-associated Calcium Signal Transducer 1 (TACSTD1); ECS-1; Epidermal Surface Antigen 1; ESA1; FLOT2; Flotillin-2; Membrane Component, Chromosome 17, Surface Marker-1 (M17S1); REG-1; Reggie-1; Reggie-2				
Human gene symbol	TACSTD1				
Entrez gene ID	4072				
SwissProt	P16422				
Unigene	542050				
Immunogen	Neuraminidase treated GLS-1 human small cell lung carcinoma cells				
Antibody target cellular localization	Plasma membrane				
Verified antibody applications	IHC (FFPE) (verified), WB (verified)				
Species reactivity	Human				
Species reactivity	Human				
Antibody application notes	Human  Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Does not react with rat, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user				
	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 1-2 ug/mL, for 30 minutes at RT, Western blot: 1-2 ug/mL, Does not react with rat, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a				
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Does not react with rat, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user				
Antibody application notes  Positive control	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 1-2 ug/mL, for 30 minutes at RT, Western blot: 1-2 ug/mL, Does not react with rat, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user				
Antibody application notes  Positive control Shipping condition	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Does not react with rat, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  HT29 cells. Breast tumor.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light,				
Antibody application notes  Positive control Shipping condition Storage Conditions	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Does not react with rat, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrale buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  HT29 cells. Breast tumor.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C				
Antibody application notes  Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Does not react with rat, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  HT29 cells. Breast tumor.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in				
Antibody application notes  Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 1-2 ug/mL, for 30 minutes at RT, Western blot: 1-2 ug/mL, poes not react with rat, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  HT29 cells. Breast tumor.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide  Guaranteed for at least 24 months from date of receipt when				
Antibody application notes  Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation Shelf life	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 1-2 ug/mL, for 30 minutes at RT, Western blot: 1-2 ug/mL, poes not react with rat, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  HT29 cells. Breast tumor.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg/mL in PBS without azide  Guaranteed for at least 24 months from date of receipt when stored as recommended				
Antibody application notes  Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation  Shelf life Cell/tissue expression	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Does not react with rat, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  HT29 cells. Breast tumor.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% SSA/0.05% szide, Purified as Pagimular in PBS without azide  Guaranteed for at least 24 months from date of receipt when stored as recommended  Epithelial cells  Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in				

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on December 15, 2025 at 03:24:36 PM. Visit product page to check for updated information before use. Product link: <a href="http://54.245.69.9/product/ep-cam-cd326-monoclonal-mouse-antibody-moc-31/">http://54.245.69.9/product/ep-cam-cd326-monoclonal-mouse-antibody-moc-31/</a>