## Mucin 1 / EMA / Episialin / CD227 Monoclonal Mouse Antibody (VU-4H5)

## **Product Description**

MAb VU-4H5 reacts with MUC1, a large transmembrane glycoprotein expressed on the ductal surface of normal glandular epithelia. The dominant epitope of MAb VU4H5 is APDTR as established with 'epitope fingerprinting'. VU-4H5 preferentially binds to under-glycosylated 'tumor' MUC1. The extracellular domain of MUC1 largely consists of a highly conserved, O-glycosylated 20 amino acids tandem repeat which can occur 30-100 times per molecule depending on the length of the allele involved. In the vast majority of human carcinomas this protein is upregulated and poorly glycosylated and appears on the cell surface in a non-polarized fashion. Antibody to EMA is useful as a pan-epithelial marker for detecting early metastatic loci of carcinoma in bone marrow or liver. Catalog number key for antibody number 0431, Anti-CA15-3|CD227|EMA|MUC1 (VU-4H5)

## Draduat attributas

Product attributes				
Antibody number	#0431			
Antibody reactivity (target)	CA15-3, CD227, EMA, MUC1			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	VU-4H5			
Isotype	IgG1, kappa			
Molecular weight	265-400 kDa			
Synonyms	Breast carcinoma-associated antigen DF3, CA15-3, Carcinoma-associated mucin Episialin, Epithelial Membrane Antigen, IZ9AG, KL-6, MAM6, MUC1-alpha, MUC1-beta, MUC1-CT, MUC1-NT, MUC1/ZD, Mucin-1 subunit beta, Peanut-reactive urinary mucin, PEM, PEMT, Polymorphic epithelial mucin, PUM, Tumor-associated epithelial membrane antigen			
Human gene symbol	MUC1			
Entrez gene ID	4582			
SwissProt	P15941			
Unigene	89603			
Immunogen	Synthetic glycosylated MUC1 60mer tandem repeat NH2-(HGVTSAPDT(GalNAc)RPAPGSTAPPAHG)3- COOH, conjugated to bovine serum albumin			
Antibody target cellular localization	Plasma membrane			
Verified antibody applications	IHC (FFPE) (verified)			
Species reactivity	Human			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user			
Antibody application notes  Positive control	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunofistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a			
	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  MCF-7 or MDA-231 cells. Breast, colon, ovarian, endometrial			
Positive control	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  MCF-7 or MDA-231 cells. Breast, colon, ovarian, endometrial carcinoma.			
Positive control Shipping condition	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  MCF-7 or MDA-231 cells. Breast, colon, ovarian, endometrial carcinoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
Positive control Shipping condition Storage Conditions	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  MCF-7 or MDA-231 cells. Breast, colon, ovarian, endometrial carcinoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C			
Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/mllion cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  MCF-7 or MDA-231 cells. Breast, colon, ovarian, endometrial carcinoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% azide, Purffied: 10.2 mg/mL in PBS/0.05% BSA/0.05% azide, Pu			
Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/mllines etls. Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  MCF-7 or MDA-231 cells. Breast, colon, ovarian, endometrial carcinoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/free: 1 mg/mL in PBS without azide  Guaranteed for at least 24 months from date of receipt when			
Positive control  Shipping condition  Storage Conditions  Regulatory status  Antibody/conjugate formulation  Shelf life	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/mllines etles, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  MCF-7 or MDA-231 cells. Breast, colon, ovarian, endometrial carcinoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/free: 1 mg/mL in PBS without azide  Guaranteed for at least 24 months from date of receipt when stored as recommended			
Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation Shelf life Cell/tissue expression	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/mllion cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  MCF-7 or MDA-231 cells. Breast, colon, ovarian, endometrial carcinoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide  Guaranteed for at least 24 months from date of receipt when stored as recommended  Epithelial cells, Mammary gland  Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in			

Call us: 800-304-5357 Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on December 15, 2025 at 03:24:30 PM. Visit product page to check for updated information before use. Product link: <a href="http://54.245.69.9/product/mucin-1-ema-episialin-cd227-monoclonal-mouse-antibody-vu-4h5/">http://54.245.69.9/product/mucin-1-ema-episialin-cd227-monoclonal-mouse-antibody-vu-4h5/</a>