Melanoma Marker Monoclonal Mouse Antibody (DT101+ BC199 + HMB45)

Product Description

This antibody cocktail recognizes two melanoma-specific proteins, which include MART-1 and gp100. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Function of gp100 is not known but it is reported to be a useful marker for melanocytes and melanomas. A cocktail of these two markers is sensitive for labeling formalin-fixed, paraffin-embedded melanomas and other tumors showing melanocytic differentiation. Catalog number key for antibody number 0703, Anti-Melanoma cells (DT101 BC199 HMB45)

Product attributes

Call us: 800-304-5357

Product attributes			
Antibody number	#0703		
Antibody reactivity (target)	Melanoma cells		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	DT101+ BC199 + HMB45		
Isotype	IgG1, kappa		
Molecular weight	20-22 kDa (doublet) (MART-1); 90-100 kDa (gp100)		
Synonyms	Melanoma antigen recognized by T-cells 1 (MART-1), MLAN-A; PMEL17		
Human gene symbol	MLANA & SILV		
Entrez gene ID	2315; 6490		
SwissProt	Q16655; P40967		
Unigene	154069 ; 95972		
Immunogen	Recombinant hMART-1 protein (DT101 & BC199); Extract of pigmented melanoma metastases from lymph nodes (HMB45)		
Antibody target cellular localization	Cytoplasmic		
Verified antibody applications	IHC (FFPE) (verified)		
	Human		
Species reactivity	Human		
Species reactivity Antibody application notes	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user		
•	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be		
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-MEL-13 and SK-MEL-19 melanoma cell lines. Human		
Antibody application notes Positive control	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/ml. for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-MEL-13 and SK-MEL-19 melanoma cell lines. Human melanoma tissues.		
Antibody application notes Positive control Shipping condition	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes to 10 minutes in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-MEL-13 and SK-MEL-19 melanoma cell lines. Human melanoma tissues. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,		
Antibody application notes Positive control Shipping condition Storage Conditions	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes to 10 minutes to 1		
Antibody application notes Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mt for 30 minutes at RT, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-MEL-13 and SK-MEL-19 melanoma cell lines. Human melanoma tissues. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified		
Antibody application notes Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 0.5-1.0 u/ml. for 30 minutes at RT, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-MEL-13 and SK-MEL-19 melanoma cell lines. Human melanoma tissues. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; DSA-free: 1 mg/mL in PBS/0.05% BSA-free: 1 mg/mL in PBS/0.05% azide, Purified; DSA-free: 1 mg/mL in PBS/0.05% azide; PURIFIED; PURIFI		
Antibody application notes Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation Shelf life	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 0.5-1.0 u/ml. for 30 minutes at RT, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-MEL-13 and SK-MEL-19 melanoma cell lines. Human melanoma tissues. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide Guaranteed for at least 24 months from date of receipt when stored as recommended		
Antibody application notes Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation Shelf life Cell/tissue expression	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mt for 30 minutes at RT, Staining of formalin-fixed dissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-MEL-13 and SK-MEL-19 melanoma cell lines. Human melanoma tissues. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA-free: 1 mg/mL in PBS without azide Guaranteed for at least 24 months from date of receipt when stored as recommended Melanocytes Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in		

Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on December 15, 2025 at 03:24:33 PM. Visit product page to check for updated information before use. Product link: http://54.245.69.9/product/melanoma-marker-monoclonal-mouse-antibody-dt101-bc199-hmb45/