CD20 / MS4A1 Monoclonal Mouse Antibody (L26)

Product Description

This antibody recognizes a protein of 30-33 kDa, which is identified as CD20. Its epitope is located in the cytoplasmic domain of CD20 and was, therefore, ascribed as CD20cy in the 5th Workshop. CD20 is a non-lg differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD20 is expressed by pre B-cells and persists during all stages of B-cell maturation but is lost upon terminal differentiation into plasma cells. This MAb can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type. Catalog number key for antibody number 0766, Anti-CD20|MS4A1 (L26)

Call us: 800-304-5357 Email: techsupport@biotium.com

Product attributes

Antibody number	#0766	
Antibody reactivity (target)	CD20, MS4A1	
Antibody type	Primary	
Host species	Mouse	
Clonality	Monoclonal	
Clone	L26	
Isotype	IgG2a, kappa	
Molecular weight	33-37 kDa	
Synonyms	APY; ATOPY; B-lymphocyte cell-surface antigen B1; Bp35; B1; Bp35; B1; B25; B25; B25; B25; B25; B25; B25; B25	
Human gene symbol	MS4A1	
Entrez gene ID	931	
SwissProt	P11836	
Unigene	712553	
Immunogen	Human tonsil B cells	
Antibody target cellular localization	Plasma membrane	
Verified antibody applications	Flow (verified), IF (verified), IHC (FFPE) (verified), WB (verified)	
Species reactivity	Human	
•		
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunofluotochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user	
•	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunofluoreschemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a	
Antibody application notes	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user	
Antibody application notes Positive control	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user Daudi, Raji and U266 cell lines. Human lymphocytes. Human lymph node or tonsil.	
Antibody application notes Positive control Shipping condition	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunofistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires bolling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user Daudi, Raji and U266 cell lines. Human lymphocytes. Human lymph node or tonsil. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,	
Antibody application notes Positive control Shipping condition Storage Conditions	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user Daudi, Raji and U266 cell lines. Human lymphocytes. Human lymph node or tonsil. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C	
Antibody application notes Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/mllion cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user Daudi, Raji and U266 cell lines. Human lymphocytes. Human lymph node or tonsil. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 9.8A fmL in PBS/0.05% BSA, Purified: 1 mg/mL in PBS/0.05% BS	
Antibody application notes Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissue seculires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user Daudi, Raji and U266 cell lines. Human lymphocytes. Human lymph node or tonsil. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide Guaranteed for at least 24 months from date of receipt when	
Antibody application notes Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation Shelf life	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user Daudi, Raji and U266 cell lines. Human lymphocytes. Human lymph node or tonsil. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg/mL in PBS without azide Guaranteed for at least 24 months from date of receipt when stored as recommended	
Antibody application notes Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation Shelf life Cell/tissue expression	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT, Western blot: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/mllion cells, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user Daudi, Raji and U266 cell lines. Human lymphocytes. Human lymph node or tonsil. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA Pare: 1 mg/mL in PBS without azide Guaranteed for at least 24 months from date of receipt when stored as recommended B-cells, Lymphocytes Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in	

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on December 15, 2025 at 10:20:22 AM. Visit product page to check for updated information before use. Product link: http://54.245.69.9/product/cd20-ms4a1-monoclonal-mouse-antibody-l26/