CD19 Monoclonal Mouse Antibody (CVID3/429)



Product Description

CD19 is a transmembrane glycoprotein that contains two extracellular immunoglobulin-like domains. CD19 is present in both benign and malignant B-cells and is considered to be the most reliable surface marker of this lineage over a wide range of maturational stages. In normal lymphoid tissue, CD19 is observed in germinal centers, in mantle zone cells, and in scattered cells of the inter-follicular areas. Anti-CD19 exhibits an overall immunoreactivity pattern similar to those of the antibodies against CD20 and CD22. However, in contrast to CD20, expression of CD19 is continuous throughout B-cell development and through terminal differentiation of B-cells into plasma cells. Anti-CD19 positivity is seen in the vast majority of B-cell neoplasms commonly at a lower intensity than normal B-cell counterparts. Plasma cell neoplasms are nearly always negative, as are T-cell neoplasms.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0429, Anti-CD19 (CVID3/429)

Product attributes

Product attributes		
Antibody number	#0429	
Antibody reactivity (target)	CD19	
Antibody type	Primary	
Host species	Mouse	
Clonality	Monoclonal	
Clone	CVID3/429	
Isotype	IgG1, kappa	
Molecular weight	95 kDa	
Synonyms	B-lymphocyte antigen CD19; B-lymphocyte surface antigen B4; CVID3; Leu-12; Lymphocyte Surface Antigen; T-cell surface antigen Leu-12	
Human gene symbol	CD19	
Entrez gene ID	930	
SwissProt	P15391	
Unigene	652262	
Immunogen	Recombinant human CD19 protein	
Antibody target cellular localization	Plasma membrane	
Verified antibody applications	Flow (surface) (verified), IF (verified)	
Species reactivity	Chimpanzee, Human, Monkey	
Species reactivity Antibody application notes	Chimpanzee, Human, Monkey Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mlL, Flow Cytometry 0.5-1 ug/mlllion cells/0.1 mL, Optimal dilution for a specific application should be determined by user	
•	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mll, Flow Cytometry 0.5-1 ug/mllion cells/0.1 ml., Optimal dilution for a	
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mll, Flow Cytometry 0.5-1 ug/mllion cells/0.1 ml., Optimal dilution for a specific application should be determined by user	
Antibody application notes Positive control	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mll, Flow Cytometry 0.5-1 ug/mllion cells/0.1 ml., Optimal dilution for a specific application should be determined by user Tonsil, lymph node, or spleen	
Antibody application notes Positive control Shipping condition	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml., Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Optimal dilution for a specific application should be determined by user Tonsil, lymph node, or spleen Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,	
Antibody application notes Positive control Shipping condition Storage Conditions	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml., Flow Cytometry 0.5-1 ug/mllion cells/0.1 ml., Optimal dilution for a specific application should be determined by user Tonsil, lymph node, or spleen Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when	
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml., Flow Cytometry 0.5-1 ug/mllion cells/0.1 ml., Optimal dilution for a specific application should be determined by user Tonsil, lymph node, or spleen Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended	
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml., Flow Cytometry 0.5-1 ug/mllion cells/0.1 ml., Optimal dilution for a specific application should be determined by user Tonsil, lymph node, or spleen Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: BSA-free 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: BSA-free 1 mg/mL in PBS/0.05% azide Purified: BSA-free 1 mg/mL in PBS/0.05% azide Purified: BSA-free 1 mg/mL in PBS/0.05% azide Purified: BSA-free 1 mg/mL in	
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml., Flow Cytometry 0.5-1 ug/mllion cells/0.1 ml., Optimal dilution for a specific application should be determined by user Tonsil, lymph node, or spleen Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg/mL in PBS without azide	
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation Antibody research areas	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mlL, Immunofluorescence: 0.5-1 ug/mlL, Flow Cytometry 0.5-1 ug/mlL individual be determined by user Tonsil, lymph node, or spleen Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.15% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS without azide Immunology Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in	

Call us: 800-304-5357 Email: btinfo@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on May 9, 2025 at 01:07:36 PM. Visit product page to check for updated information before use. Product link: http://54.245.69.9/product/monoclonal-anti-cd19-cvid3429/