CD2 Monoclonal Mouse Antibody (UMCD2)

Product Description

CD2 interacts through its amino-terminal domain with the extracellular domain of CD58 (also designated CD2 ligand) to mediate cell adhesion. CD2/CD58 binding can enhance antigen-specific T cell activation. CD2 is a transmembrane glycoprotein that is expressed on peripheral blood T lymphocytes, NK cells and thymocytes. CD58 is a heavily glycosylated protein with a broad tissue distribution in hematopoietic and other cells, including endothelium. Interaction between CD2 and its counter receptor LFA3 (CD58) on opposing cells optimizes immune system recognition, thereby facilitating communication between helper T lymphocytes and antigen-presenting cells, as well as between cytolytic effectors and target cells.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 1235, Anti-CD2 (UMCD2)

Product attributes

Product attributes			
Antibody number	#1235		
Antibody reactivity (target)	CD2		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	UMCD2		
Isotype	IgG2a, kappa		
Molecular weight	50 kDa		
Synonyms	Erythrocyte receptor; LFA-2; LFA-3 receptor; Ly-37; Lymphocyte function antigen 2; Rosette receptor; SRBC; T-cell surface antigen CD2; T-cell surface antigen T11/Leu-5; T11		
Human gene symbol	CD2		
Entrez gene ID	914		
SwissProt	P06729		
Unigene	523500		
Immunogen	Human Thymocytes/Sezary T cells		
Antibody target cellular localization	Plasma membrane		
Verified antibody applications	Flow (verified), WB (verified)		
Species reactivity	Human		
Expected antibody applications	Flow, surface (published for clone), Functional studies (published for clone), WB (published for clone)		
applications	(published for clone), WB (published for clone) Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml., Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Optimal dilution for a		
applications Antibody application notes	(published for clone), WB (published for clone) Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
applications Antibody application notes Positive control	(published for clone), WB (published for clone) Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immonfluorescence: 0.5-1 ug/mltibody. Brown Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user 293T or Jurkat cells. Lymph node and tonsil		
applications Antibody application notes Positive control Shipping condition	(published for clone), WB (published for clone) Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user 293T or Jurkat cells. Lymph node and tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,		
applications Antibody application notes Positive control Shipping condition Storage Conditions	(published for clone), WB (published for clone) Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user 293T or Jurkat cells. Lymph node and tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when		
applications Antibody application notes Positive control Shipping condition Storage Conditions Shelf life	(published for clone), WB (published for clone) Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user 293T or Jurkat cells. Lymph node and tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended		
applications Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	(published for clone), WB (published for clone) Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user 293T or Jurkat cells. Lymph node and tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified; 0.2 mg/mL in PBS/0.05% BSA, Purified; 0.2 mg/mL in PBS/0.05% BSA, Purified; 0.2 mg/mL in PBS/0.05% BSA, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% aside, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% aside, Purified; BSA-free: 1 mg/mL in		
applications Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	(published for clone), WB (published for clone) Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user 293T or Jurkat cells. Lymph node and tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purffled, BSA-free: 1 mg/mL in PBS without azide		

Call us: 800-304-5357 Email: btinfo@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

References

Note: References for this clone sold by other suppliers may be listed for expected applications.

Cellular Immunol (1993) 150(2): 235-246. (Flow, surface: WB: functional studies)

This datasheet was generated on May 9, 2025 at 03:06:57 AM. Visit product page to check for updated information before use. Product link: http://54.245.69.9/product/monoclonal-anti-cd2-umcd2/